

病名：胆道癌

レジメン名：GS療法

1コース期間(休薬期間含む) 21 日

薬剤名/ルート・用法	基準投与量	割合	1/1	1/2	1/3	1/4	1/5	1/6	1/7	1/8	1/9	1/10	1/11	1/12	1/13	1/14	1/15	1/16	1/17	1/18	1/19	1/20	1/21
			火	水	木	金	土	日	月	火	水	木	金	土	日	月	火	水	木	金	土	日	月
			day1 未適用	day2~day7 休薬							day8 未適用	day9~day21 休薬											
注射																							
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)加算A	00:01		未適用																				
パロチロン点滴静注バッグ® 0.75mg 50ml	1 袋																						
デキサート注射液6.6mg 2mL	1 V																						
末梢点滴ケモ(持続)	00:05		未適用																				
生理食塩液「ヒカリ」100mL	1 V																						
●ゲムシタビ点滴静注 200mg	1000 mg/m <sup>2</sup>	100																					
末梢点滴ケモ CSTD(側管)	00:10		未適用																				
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)	00:15		未適用																				
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)加算A	00:01									未適用													
パロチロン点滴静注バッグ® 0.75mg 50ml	1 袋																						
デキサート注射液6.6mg 2mL	1 V																						
末梢点滴ケモ(持続)	00:05									未適用													
生理食塩液「ヒカリ」100mL	1 V																						
●ゲムシタビ点滴静注 200mg	1000 mg/m <sup>2</sup>	100																					
末梢点滴ケモ CSTD(側管)	00:10									未適用													
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)	00:15									未適用													
処方																							
注意コメント	S-1:day1夕~15朝 60~100mg/day																						