

病名:前立腺癌

レジメン名:低用量ED(エストラムスチン・ドセタキセル)療法

1コース期間(休薬期間含む) 21 日

薬剤名/ルート・用法	基準投与量	割合	1/1	1/2	1/3	1/4	1/5	1/6	1/7	1/8	1/9	1/10	1/11	1/12	1/13	1/14	1/15	1/16	1/17	1/18	1/19	1/20	1/21
			火	水	木	金	土	日	月	火	水	木	金	土	日	月	火	水	木	金	土	日	月
			day1 休薬	day2 未適用	day3~day8 休薬						day9 未適用	day10~day21 休薬											
注射																							
生理食塩液「ヒカリ」100mL	1 V																						
デキサト注射液6.6mg 2mL	1 V																						
ガスター注射液 20mg 2mL	1 A																						
末梢点滴ケモ(持続)	00:05																						
ブドウ糖注 5% 250mL	1 V																						
●ドセタキセル注 ☆20mg	25 mg/m ²	100																					
末梢点滴ケモ CSTD(側管)	00:10																						
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)	00:15																						
生理食塩液「ヒカリ」100mL	1 V																						
デキサト注射液6.6mg 2mL	1 V																						
ガスター注射液 20mg 2mL	1 A																						
末梢点滴ケモ(持続)	00:05																						
ブドウ糖注 5% 250mL	1 V																						
●ドセタキセル注 ☆20mg	25 mg/m ²	100																					
末梢点滴ケモ CSTD(側管)	00:10																						
生理食塩液「ヒカリ」50mL	1 V																						
末梢点滴ケモ(持続)	00:15																						
処方																							
注意コメント	day1-3、day8-10 朝夕 エストラサイト 4cap day1-21 朝 デカロン0.5mg 2T																						